



FRUIT-FLIES (DIPTERA: TEPHRITIDAE) FROM NORTH-EAST INDIA

KAUSHIK KR. BHATTACHARYA, P. PARUI, DHIRITI BANERJEE*

Zoological Survey of India, Kolkata-700053.

*E-mail : dhritybanerjee@gmail.com

INTRODUCTION

Northeast India consists the present day Assam or Assam valley and rest of the contiguous - Arunachal Pradesh, Meghalaya, Mizoram, Nagaland and former princely states of Manipur and Tripura, state of Sikkim and parts of North Bengal (districts of Darjeeling, Jalpaiguri and Koch Bihar). The present paper based on old collections present in Zoological Survey of India records 6 species under 4 genera from North-East India.

Order DIPTERA

Family TEPHRITIDAE

Subfamily DACINAE

Tribe DACINI

Genus *Bactrocera* Macquart

1835. *Bactrocera* Macquart, *Hist. nat. Ins. Dipt.*, 2: 452,453. Type-species: *Bactrocera longicornis* Macquart.

Subgenus *Zeugodacus* Hendel

1927. *Zeugodacus* Hendel, in Lindner, *Fliegen palaearkt. Reg.*, 49: 26. Type-species: *Bactrocera caudatus* Fabricius

Key to the species

1. Costal band interrupted and very thin, abdomen largely black.....*Bactrocera (Zeugodacus) scutellaris* (Bezzi). (Fig. a, Plate-1).
- Costal band distinct and largely enlarged at apex; abdomen largely yellow..... *Bactrocera (Zeugodacus) tau* (Walker) (Fig. b, Plate-1).

Bactrocera (Zeugodacus) scutellaris (Bezzi)

1913. *Bactrocera scutellaris* Bezzi, *Mem. Indian Mus.*, 3: 98. Type-loc: India: (Meghalaya: Shillong; West

Bengal: Suiliguri, Kurseong; Uttarakhand: Kumaon, Bhowali).

Material examined: 1 , Dentam, West Sikkim dist., 18.x.1995, coll. S.C. Mitra.

Distribution: India: Sikkim, Arunachal Pradesh, Bihar, Himachal Pradesh, Karnataka, Meghalaya, Uttarakhand, West Bengal. *Elsewhere:* China, Myanmar, Nepal, Pakistan, Sri Lanka, Thailand.

Remarks: This species is being reported for the first time from Sikkim.

Subgenus *Zeugodacus* Hendel

1927. *Zeugodacus* Hendel, in Lindner, *Fliegen palaearkt. Reg.*, 49: 26. Type-species, *Dacus caudatus* Fabricius.

Bactrocera (Zeugodacus) tau (Walker)

1849. *Dasyneura tau* Walker, *List Dipt. colln Br. Mus.*, 4: 1074. Type-Loc: Foochow, China.

Material examined: 1 , Tuikual south, Aizawl dist., 31.iii.94, coll. A.K. Hazra; 1 , Belonia, South Tripura dist., 25.iii.91, coll. G.K. Srivastava.

Distribution: India: Mizoram, Tripura, Andaman & Nicobar Islands, Arunachal Pradesh, Bihar, Haryana, Himachal Pradesh, Jammu & Kashmir, Karnataka, Maharashtra, Meghalaya, Punjab, Sikkim, Tamil Nadu, Uttarakhand, West Bengal. *Elsewhere:* Bhutan, Myanmar, China, Taiwan, Laos, Thailand, Cambodia, Vietnam, Malaysia, Singapore, Brunei, Indonesia.

Remarks: This species is being reported for the first time from Mizoram and Tripura.

Subfamily TEPHRITINAE

Tribe TEPHRELLINI

Subtribe TEPHRELLINA

Genus *Oxyaciura* Hendel

1927. *Oxyaciura* Hendel, In Lindner (ed.) *Die Fliegen der Palaearktischen Region*, 5: 111. Type species, *Aciura tibialis* Robineau-Desvoidy.

Oxyaciura xanthotricha (Bezzi)

1913. *Aciura xanthotricha* Bezzi, *Mem. Indian Mus.*, 3: 151. Type-loc: India: Uttarakhand (Dhikata).

Material examined: 1 ♂, Singhat, Churachandpur dist., 18.xii.92, coll. R.S. Mridha; 1 ♂, 2 ♀, Pangian Village, Churachandpur dist., 17.xii.92, coll. R.S. Mridha; 1 ♂, Malsa, Moreh, Chandel dist., 25.xii.92, coll. R.S. Mridha; 2 ♀, Rangpo, East Sikkim dist., 12.x.95, coll. S.C. Mitra.

Diagnosis: The species has flat, white, occipital setae (bristles) and postocellar bristles and scapular hairs lacking, typical of Tephritinae, but lacks recumbent, gray white setae over mesonotum and the 6th abdominal segment of the female is distinctly shorter than the 5th as in Trypetinae; upper occipital setae are flat and white; with only 3 hyaline marks in hind margin of wing; basal segment of ovipositor approximately as long as remainder of abdomen; piercer slender, tapered to a long thin point, extended ovipositor. (Fig. c, Plate-1).

Distribution: India: Manipur, Sikkim, Himachal Pradesh, Meghalaya, Uttarakhand. *Elsewhere*: Indonesia, Myanmar, Sri Lanka, Thailand, Vietnam, China.

Remarks: This species is being reported for the first time from Manipur and Sikkim.

Tribe TEPHRITINI

Genus *Campiglossa* Rondani

1870. *Campiglossa* Rondani, *Bull. Soc. Entomol. Ital.*, 2: 121. Type-species, *Tephritis irrorata* Fallen.

Key to the species

1. Wing entirely dark brown with numerous dark brown spots along periphery, cell r4+5 with a hyaline spot at apex..... *Campiglossa cribellata* Bezzi. (Fig.d, Plate-2).

- Posterior region of wing more hyaline with few dark patches or spots, cell r4+5 with an apical hyaline spot.....*Campiglossa gemma* (Hering) (Fig.e, Plate-2).

Campiglossa cribellata Bezzi

1913. *Campiglossa cribellata* Bezzi, *Mem. Indian Mus.*, 3: 161. Type-loc: India (West Bengal: Kurseong).

Material examined: 4 ♂, Nongpoh, Ri-Bhoi dist., 22.xii.1978, coll. P.T. Cherian.

Distribution: India: Meghalaya, Bihar, Haryana, Himachal Pradesh, Sikkim, West Bengal. *Elsewhere*: Nepal.

Remarks: This species is being reported for the first time from Meghalaya.

Campiglossa gemma (Hering)

1939. *Paroxyna gemma* Hering, *Verh. VII Intern. Kongr. Entomol.*, 1: 183. Type-loc: India: Tamil Nadu (Kodaikanal).

Material examined: 6 ♂, 2 ♀, Far Pak, Blue Mountain, Chhimpuitui dist., 9.iv.1994, coll. A.K. Hazra.

Distribution: India: Mizoram, Jammu & Kashmir (Ladakh), Karnataka, Tamil Nadu. *Elsewhere*: Nil.

Remarks: This species is being reported for the first time from Mizoram and is endemic to India.

Genus *Dioxyina* Frey

1954. *Dioxyina* Frey, *Commentat. Biol. Soc. Sci. Fenn.* (1944), 8(10): 62. Type-species, *Trypeta sororcula* Wiedemann.

Dioxyina sororcula (Wiedemann)

1830. *Trypeta sororcula* Wiedemann, *Aussereurop. Zweifl. Insekt.*, 2: 509. Type-loc: Teneriffe, Canary Islands.

Material examined: 1 ♂, Far Pak, Blue Mountain, Chhimpuitui dist., 9.iv.1994, coll. A.K. Hazra; 1 ♀, Belonia, South Tripura dist., 25.iii.91, coll. G.K. Srivastava.

Diagnosis: Head longer than high, epistoma and sides of face protruded, proboscis slender, elongate, geniculate, two pairs of inferior and superior fronto-orbital bristles present, dorsocentral bristles situated distinctly in front of

a line between the anterior supraalars; subcostal cell of wing dark brown; base of ovipositor shining black, subequal in length to terga 4-6. (Fig. f, Plate-2).

Distribution: India: Mizoram, Tripura, Andhra Pradesh, Assam, Bihar, Chandigarh, Haryana, Himachal Pradesh, Jammu & Kashmir, Jharkhand, Karnataka, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Nagaland, Orissa, Punjab, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal. *Elsewhere:* Widespread over Tropics and subtropics of the world, Japan, Australia, S. Europe, Korea, Southern Canada to Chile. Belonia, South Tripura dist., 25.iii.91, coll. G.K. Srivastava.

Diagnosis: Head longer than high, epistoma and sides of face protruded, proboscis slender, elongate, geniculate, two pairs of inferior and superior fronto-orbital bristles present, dorsocentral bristles situated distinctly in front of a line between the anterior supraalars; subcostal cell of wing dark brown; base of ovipositor

shinning black, subequal in length to terga 4-6. (Fig. f, Plate-2).

Distribution: India: Mizoram, Tripura, Andhra Pradesh, Assam, Bihar, Chandigarh, Haryana, Himachal Pradesh, Jammu & Kashmir, Jharkhand, Karnataka, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Nagaland, Orissa, Punjab, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal. *Elsewhere:* Widespread over Tropics and subtropics of the world, Japan, Australia, S. Europe, Korea, Southern Canada to Chile.

Remarks: This species is being reported for the first time from Mizoram and Tripura.

ACKNOWLEDGEMENTS

The authors are indebted to Dr. K. Venkataraman, Director, Zoological Survey of India, for his kind support and help and Dr. Animesh Bal, Scientist-'F' for his constant encouragement. The authors express their thanks to Shri Sudipta Pande, Laboratory Assistant, for assisting in preparing the plates.

REFERENCES

- Agarwal, M.L and Sueyoshi, M. 2005. Catalogue of Indian Fruit flies (Diptera: Tephritidae). *Oriental Insects*, **39**: 371-433.
- Bezzi, M. 1913. Indian trypanoids (fruit-flies) in the collection of the Indian Museum, Calcutta. *Memoirs of Indian Museum*, **3**: 53-175.
- Bezzi, M. 1915. Two new species of fruit flies from southern India. *Bulletin of Entomological Research*, **5**: 153-154.
- Bezzi, M. 1916. On the fruit-flies of the genus *Dacus* (s.l.) occurring in India, Burma, and Ceylon. *Bulletin of Entomological Research*, **7**: 99-121.
- Bhattacharya, K.K., Parui, P., Banerjee, D. 2012. *Sphaeniscus quadrincisus* (Wiedemann, 1824) (Diptera:Tephritidae), a new record from Himachal Pradesh. *Records of Zoological Survey of India*, (in press).
- Brunetti, E. 1917. Diptera of Simla District. *Rec. Ind. Mus.*, **13**: 59-101.
- David, K.J. and Ramani, S. 2011. An Illustrated Key to Fruit flies (Diptera:Tephritidae) from Peninsular India and Andaman & Nicobar Islands, *Zootaxa*, **3021**: 1-31.
- Drew, R.A.I., Hancock, D.L. and White, I.M. (1998) Revision of the tropical fruit flies (Diptera: Tephritidae: Dacinae) of Southeast Asia. II. *Dacus* Fabricius. *Invertebrate Taxonomy*, **12**: 567-654.
- Drew, R.A.I and Raghu, S. 2002. The fruit fly fauna (Diptera : Tephritidae: Dacinae) of the rain forest habitat of the Western Ghats, India. *The Raffles Bulletin of Zoology*, **50** (2): 327-352.

- Evenhuis, N.L., Pape, T., Pont, A.C. and Thompson, F.C. 2008. *Biosystematic Database of World Diptera, Version 10*, <http://www.diptera.org/biosys.htm>.
- Hancock, D.L. and Drew, R.A.I. 2005. New genera, species and records of Adramini (Diptera : Tephritidae : Trypetinae) from the South Pacific and southern Asia. *Australian Entomologist*, **32** (1): 5-16.
- Hardy, D.E. 1971. Diptera: Tephritidae from Ceylon. *Entomologica Scandinavica. Supplement*, **1**: 287-292.
- Hardy, D.E. 1973. The fruit flies (Tephritidae : Diptera) of Thailand and bordering countries. *Pacific Insects Monograph*, **31**:1-353.
- Hardy, D.E. 1974. The fruit flies of the Philippines (Diptera : Tephritidae). *Pacific Insects Monograph*, **32**: 1-266.
- Hering, E.M. 1938. Entomological results from the Swedish Expedition to Burma and British India. Diptera: Fam. Trypetidae. [23. Beitrag zur Kenntnis der Trypetidae]. *Arkive for Zoologi*, **30A** (25): 56 pp.
- Hering, E.M. 1941. Entomological results from the Swedish Expedition to Burma and British India. Diptera: Trypetidae. Nachtrag [30. Beitrag zur Kenntnis der Trypetidae]. *Arkive for Zoologi*, **33B** (11): 7 pp.
- Hering, E.M. 1956. Trypetidae (Dipt.) von Ceylon (53. Beitrag zur Kenntnis der Trypetidae). *Verhandlungen der Naturforschenden Gesellschaft in Basel*, **67**: 62-74.
- Kapoor, V.C. 1971. Four new species of fruit flies (Tephritidae) in India. *Oriental Insects*, **5**: 477-482.
- Kapoor, V.C. 1993. *Indian fruit flies (Insecta: Diptera: Tephritidae)*. Oxford & IBH, New Delhi, 228 pp.
- Kapoor, V.C., Hardy, D.E., Agarwal, M.L. and Grewal, J.S. 1980. *Fruit fly (Diptera: Tephritidae) systematics of the Indian subcontinent*. Export India, Jullundur, India: 113 pp.
- Korneyev, V.A. 1999. Phylogenetic Relationships among Higher Groups of Tephritidae. In: Aluja, M & Norrbom, A.L. (Eds.), *Fruit Flies (Tephritidae): Phylogeny and Evolution of Behaviour*. CRC Press, Boca Raton, pp 73-113.
- Munro, H.K. 1935. Records of Indian Trypetidae (Diptera) with descriptions of some apparently new species. *Records of Indian Museum*, **37**: 15-27.
- Munro, H.K. 1938. Studies on Indian Trypetidae (Diptera). *Records of Indian Museum*, **40**: 21-37.
- Munro, H.K. 1939. The fruit fly, *Dacus ferrugineus* Fabr., and its variety *dorsalis* Hendel in north west India. *Indian Journal of Entomology*, **1**: 101-105.
- Pape, T. and Thomson, F.C. (eds) 2012. *Systema Dipteroorum* (version 2.0, Jan 2011). In: Species 2000 & IT IS Catalogue of Life, 2012 Annual Checklist (Bisby F., Roskov Y., Culham A., Orrell T., Nicolson D., Paglinawan L., Bailley N., Appeltans W., Kirk P., Bourgoin T., Baillargeon G., Ouvrard D., eds).
- Perkins, F.A. 1938. Studies in Oriental and Australian Trypanidae-Part II. Adraminae and Dacinae from India, Ceylon, Malaya, Sumatra, Java, Borneo, Philippine Islands, and Formosa. *Proceedings of Royal Society of Queensland*, **49**: 120-144.
- Radhakrishnan, C. 1999. A new species of *Bactrocera* Macquart (Diptera: Tephritidae: Dacinae) from southern India. *Rec. zool. Surv. India*, **97**(4): 1-4.
- Senior-White, R.A. 1922. Notes on Indian Diptera 1. Diptera from the Khasia Hills. 2. Tabanidae in the collection of the forest zoologist. 3. New species of Diptera from the Indian Region. *Memoirs of the Department of Agriculture in India Entomological Series*, **7**: 83-170.

Senior-White, R.A. 1924. *Catalogue of Indian insects. Part 4-Trypetidae (Trypaneidae)*. Government of India, Central Publication Branch, Calcutta, 33 pp.


White, I.M. and Hancock, D.L. 1997. *CABIKEY to the Dacini (Diptera, Tephritidae) of the Asia-Pacific-Australasian Regions*. CABINTERNATIONAL, Wallingford. Windows CD.

White, I.M. and Elson-Harris, M.M. 1992. *Fruit-flies of economic significance: Their identification and Bionomics*. CAB Int., Wallingford, 601 pages.

Plate -1



a. *Bactrocera (Zeugodacus) scutellaris* (Bezzi)



a. wing




b. *Bactrocera (Zeugodacus) tau* (Walker)



b. wing



c. *Oxyaciura xanthotricha* (Bezzi)



c. wing

Plate - 2



d. *Campiglossa cribellata* Bezzi



d. wing



e. *Campiglossa gemma* (Hering)



e. wing



f. *Dioxyna sororcula* (Wiedemann)



f. wing