

IX ARACHNIDA, I.

A. CHELIFERA.

By E. ELLIGSEN.

Chelifer javanus, Thorell.

2♂, 1♀ Kobo, 400 ft., xi—xii-1911, under logs. S. W Kemp, leg.

The species was originally described from Java, and was afterwards recorded from several other localities in the Oriental region, such as Burma, Kelantan (Malay Peninsula), Bismarck Archipelago, and the Marshall Islands.

Chelifer superbus, With.

1♀ Kobo, 400 ft., xi—xii-1911, under logs. S. W Kemp, leg.

Only known from the Celebes, from where Mr. With obtained the types.

B. PEDIPALPI.

By F. H. GRAVELY, M.Sc., Assistant Superintendent,
Indian Museum.

Uroproctus assamensis, Stoliczka.

J.A.S.B., xxxviii (2).

Kobo, 400 ft., 3—7-xii-11, under logs. Banks of Siyom River below Damda, 1300 ft., 1-ii-12. Beside stream below Balek, 26-iii-12. Rotung, 1300 ft., 21 and 25-xii-11 and 12-iii-12, under logs and under stones. Upper Rotung, ca. 2000 ft., 1—10-i-12.

This species is the only *Uroproctus* yet known, and appears to be abundant throughout the foot-hills of the E. Himalayas and in many parts of Assam. In Burma it is replaced by the genus *Hypoctonus*, of which genus one species was described not long ago from Sylhet in W. Assam, and another still more recently from the foot of the Himalayas in the Darjeeling district. The western end of Assam and the E. Himalayas at present appear therefore to have a greater affinity to the Burmese (but not Malaysian) Thelyphonid fauna than does the country east of them; but many